

# ZEST – Zocular Eyelid System Treatment – improves contact lens discomfort caused by MGD


Srihari Narayanan, Erika Bendert, Kathryn Owen, Thelzy Zamaron, Christian Lopez, Narges Kasraie Rosenberg School of Optometry, University of the Incarnate Word, San Antonio, TX, United States


### Introduction

- Contact lens discomfort (CLD) reduces CL tolerance and wear times. Meibomian Gland Dysfunction (MGD) often causes CLD.
- We have previously shown that lid margin debridement-scaling improves MGD signs and decreases inflammatory marker matrix metalloproteinase-9 (MMP-9)
- Zocular is a novel "okra-based" formulation to enhance lid margin hygiene. The Zocular Eyelid System Treatment (ZEST) is performed in-office by the clinician
- The purpose of this clinical trial (NCT03968731) was to study improvement in CLD associated with MGD using the ZEST protocol.

### Methods

- Thirty CLD patients with MGD (8 men; 26.5 +/- 7.51 years) consented to the study; ZEST was performed
  in all eyelids on the first visit
- Patients reported number of hours of CL wear per day and completed the CLDEQ-8 questionnaire prior to treatment and at 2-weeks and 1-month post treatment
- Quidel Inflammadry MMP-9 test was done prior to and 1-month post-treatment; Non-invasive tear break-up (NITBUT) was measured at each visit; Data was analyzed by paired samples t-test.









## Results


Decreased Lid Wiper epitheliopathy




### **Decreased CLDEQ-8 scores Post ZEST Tx**




**Decreased OSDI scores Post ZEST Tx** 



**Increased CL Wear Time Post ZEST Tx** 



**Decreased MMP-9 Post ZEST Tx** 



### Conclusions

- Results show significant increase in CL-wear time and betterment in CLD symptoms following ZEST treatment
- The inflammatory marker MMP-9 was significantly decreased in the group post-treatment
- NITBUT showed a trend of betterment at the conclusion of the study
- The outcomes suggest that improvement in lid margin hygiene subsequent to ZEST protocol contributes to better comfort and tolerance of contact lenses as well as an ocular surface environment with decreased inflammation

## Acknowledgements

#### Funding:

UIW Faculty Development Research Grant UIW Summer Fellowship Training Program Zocular – Provided ZEST Kits